

ULTRA-LUMINOUS X-RAY SOURCES: EVIDENCE FOR VERY EFFICIENT FORMATION OF POPULATION III STARS CONTRIBUTING TO THE COSMIC NEAR-INFRARED BACKGROUND EXCESS?

HARUKA MII AND TOMONORI TOTANI
Department of Astronomy, Kyoto University
Sakyo-ku, Kyoto 606-8502, Japan
Draft version February 2, 2008

ABSTRACT

Accumulating evidence indicates that some of ultra-luminous X-ray sources (ULXs) are intermediate mass black holes (IMBHs), but the formation process of IMBHs is unknown. One possibility is that they were formed as remnants of population III (Pop III) stars, but it has been thought that the probability of being an ULX is too low for IMBHs distributed in galactic haloes to account for the observed number of ULXs. Here we show that the number of ULXs can be explained by such halo IMBHs passing through a dense molecular cloud, if Pop III star formation is very efficient as recently suggested by the excess of the cosmic near-infrared background radiation that cannot be accounted for by normal galaxy populations. We calculate the luminosity function of X-ray sources in our scenario and find that it is consistent with observed data. Our scenario can explain that ULXs are preferentially found at outskirts of large gas concentrations in star forming regions. A few important physical effects are pointed out and discussed, including gas dynamical friction, radiative efficiency of accretion flow, and radiative feedback to ambient medium. ULXs could last for $\sim 10^{5-6}$ yr to emit a total energy of $\sim 10^{53}$ erg, which is sufficient to power the ionized expanding nebulae found by optical observations. *Subject headings:* black hole physics — X-rays: galaxies — infrared:general

1. INTRODUCTION

Ultra-Luminous X-ray sources (ULXs, Makishima et al. 2000 and references therein) are bright X-ray sources having luminosities greater than $\sim 3 \times 10^{39}$ ergs s⁻¹ found in off-nuclear regions of nearby galaxies. The luminosity exceeds the Eddington limit of a $\sim 20M_{\odot}$ black hole (BH) that is the maximum mass expected from normal stellar evolutionary paths (Fryer & Kalogera 2001), and their origin is now a matter of hot debate. A few models have been proposed to explain the high luminosity from ordinary stellar mass black holes, such as geometrical beaming (King et al. 2001; King 2004) or super-Eddington accretion (Begelman 2002). However, clear observational evidence for these has not yet been found. On the other hand, the high luminosity may also be explained by intermediate mass black holes (IMBHs) of $M_{BH} \gtrsim 100M_{\odot}$, without violating the Eddington limit. Recently evidence for IMBHs has been accumulating for many ULXs. They show lower temperature (~ 0.1 keV) blackbody components than that of the Galactic black hole binaries, which is consistent with IMBHs being ULXs (Miller et al. 2004). It is known that, for X-ray binaries and active galactic nuclei (AGNs), the break frequency of the power density spectrum of X-ray flux variability scales with the black hole mass, and the break frequency of ULXs is indeed between those of black hole binaries and AGNs, again indicating intermediate BH mass for ULXs (Cropper et al. 2004). It appears that at least a part of ULXs, especially the most luminous ones, are IMBHs. (See e.g., Colbert & Miller 2004; Mushotzky 2004 for recent reviews.)

However, the formation of IMBHs to become ULXs is a challenging problem. One possibility is formation in dense stellar regions or stellar clusters (Portegies Zwart & McMillan 2002). However, though ULXs are preferentially found in actively star forming regions, their position is near but not coincident with such stellar clusters, disfavoring this scenario

(Zezas & Fabbiano 2002).

On the other hand, it is theoretically expected that IMBHs form as remnants of massive population III (Pop III) stars (e.g., Schneider et al. 2002), and they may become ULXs if they accrete sufficient gas in nearby galaxies (Madau & Rees 2001). However, it is not easy to explain the observed number of ULXs by this scenario. Islam, Taylor & Silk (2004) considered a possibility that free-floating IMBHs accrete from a residual core of gas associated with the IMBHs from their formation epoch. It is highly uncertain whether such gas core can survive until present from the early universe, and this scenario cannot explain the majority of ULXs that are associated with star forming regions. Volonteri & Perna (2005) recently performed a similar calculation, and they found even more pessimistic prediction for the observable ULX number. Krolik (2004) considered a possibility that free-floating IMBHs become ULXs when they enter into dense molecular clouds in star forming regions of galactic disks, so that the association with star forming regions can be explained. Krolik considered a hypothetical “disk” population of Pop III stars that form within a galactic disk, from primordial gas that has somehow been preserved until the disk formation. This is because the number of ULXs would be too small if, as normally expected, Pop III remnants are distributed in a galactic halo. It is an interesting possibility, but it is difficult to quantitatively estimate how much amount of such gas can survive during the formation of galactic disks.

The efficiency of Pop III star formation is highly uncertain, but there are some observational hints. Independent groups (Wright & Reese 2000; Cambr sy et al. 2001; Matsumoto et al. 2004) reported detections of the cosmic near-infrared background radiation (CNIB), which cannot be explained by normal galaxy populations (Totani et al. 2001). The excess is particularly large at $\sim 1.5\mu\text{m}$, which is at $\sim 5-6\sigma$ level according to the IRTS data of Matsumoto et al., though there may still be large systematic uncertainty in the CNIB measurements, such as zodiacal light subtraction. It has been

shown that efficient formation of Pop III stars before $z \sim 8$ can explain this unaccounted excess of CNIB (Santos, Bromm, & Kamionkowski 2002; Salvaterra & Ferrara 2003); the spectral peak of the excess at $\sim 1.5\mu\text{m}$ corresponds to the redshifted Lyman α line emission.

The required amount of Pop III stars to explain the CNIB is somewhat extreme (see, e.g., Madau & Silk 2005); about $\sim 10\%$ of the cosmic baryons must be converted into Pop III stars. However, paucity of theoretical understanding of Pop III stars does not allow to reject this possibility completely, and it is the only one realistic explanation of the CNIB excess, except for the systematic uncertainty in the CNIB measurements. Theoretical studies on Pop III star formation indicate that they are likely more massive than normal stars, and most of Pop III stars could be massive enough to collapse into IMBHs without any metal ejection, meeting the metallicity constraints from quasar absorption line systems. The consequence of this scenario is that there should be IMBHs with $M_{\text{BH}} \sim 10^2\text{--}10^3 M_\odot$ in halos of nearby galaxies, whose mass fraction in the baryonic matter is $f_{\text{III}} \equiv \Omega_{\text{IMBH}}/\Omega_B \sim 0.1$, where $\Omega_B \sim 0.05$ is the cosmic baryon density in the standard definition. This is consistent with observational constraints from gravitational lensing experiments (Wambsganss 2002) and from stellar dynamics such as heating of disk stars (Wasserman & Salpeter 1994), disruption of globular clusters (Murali et al. 2000), and disruption of halo wide binaries (Yoo et al. 2004).

In this paper we show that, if this is the case, such IMBHs passing through dense molecular clouds of star forming regions should have a significant contribution to the observed ULX population, especially to the highest luminosity ones. In §2, we give an order-of-magnitude estimate of the expected ULX number. We discuss two relevant physical processes in §3 (gas dynamical friction) and §4 (radiative efficiency of accretion disks). We will present a more detailed calculation of the predicted ULX luminosity function in §5. Radiative feedback and the ULX duty cycle will be discussed in §6. Finally some discussions, conclusions, and predictions will be presented in §7.

2. NUMBER OF ULXS: AN ORDER-OF-MAGNITUDE ANALYSIS

Detectability of isolated BHs accreting interstellar gas has been studied in a number of publications (Fujita et al. 1998, Agol & Kamionkowski 2002 and references therein), and we follow these for the basic formulations. As a typical galaxy, we choose parameters of our Galaxy with a total halo mass of $M_{\text{halo}} = 10^{12} M_\odot$ and circular velocity of $V_c = 220 \text{ km s}^{-1}$. Luminosity of a black hole with mass M_{BH} expected from the Bondi-Hoyle accretion in a gas cloud is given by

$$L_X(n, v) = \dot{m} c^2 \eta = \frac{4\pi\eta c^2 G^2 M_{\text{BH}}^2 \rho}{(v^2 + \sigma_{\text{MC}}^2 + c_s^2)^{3/2}} \\ = 4 \times 10^{37} \eta_{-1} M_2^2 n_2 \tilde{v}_1^{-3} \text{ ergs s}^{-1}, \quad (1)$$

where $\dot{m}$ is the gas mass accretion rate, v the IMBH velocity relative to the gas cloud, and ρ , σ_{MC} , and c_s are the density, the turbulent velocity, and the sound speed of a molecular cloud, respectively. In the third equation, we used $M_2 \equiv M_{\text{BH}}/(100 M_\odot)$, $n_2 \equiv n/(100 \text{ cm}^{-3})$, and $\tilde{v}_1 \equiv \tilde{v}/(10 \text{ km s}^{-1})$, where $n = \rho/(\mu m_p)$ is the hydrogen number density, m_p the proton mass, $\mu = 0.13$ the mean molecular weight, and $\tilde{v} \equiv (v^2 + \sigma_{\text{MC}}^2 + c_s^2)^{1/2}$. The parameter $\eta \equiv 0.1\eta_{-1}$ represents not only the radiative efficiency of accreting gas but also includes the uncertainty in the Bondi-Hoyle formula of accretion rate. The turbulent velocity σ_{MC} scales with density as

$\sigma_{\text{MC}} \simeq 3.7 n_2^{-0.35} \text{ km s}^{-1}$ (Larson 1981). The sound speed c_s depends on the gas temperature, which is $c_s \sim 0.3 \text{ km s}^{-1}$ for cold molecular gas ($T = 10 \text{ K}$) but $\sim 10 \text{ km s}^{-1}$ for ionized hot gas ($T \sim 10^4 \text{ K}$).

The number of ULXs in the Galaxy, whose luminosity and ambient gas density are L_X and n , respectively, is then given as:

$$\frac{\partial^2 N_{\text{ULX}}}{\partial n \partial L_X} = N_{\text{IMBH}} f_{\text{disk}} \frac{d f_n(n)}{dn} \frac{d f_v(v)}{dv} \left| \frac{\partial v(n, L_X)}{\partial L_X} \right|, \quad (2)$$

where $N_{\text{IMBH}} = f_{\text{III}}(\Omega_B/\Omega_M)M_{\text{halo}}/M_{\text{BH}}$ is the total number of IMBHs in our Galactic halo, the cosmological density parameter $\Omega_M = 0.3$, f_{disk} the fraction of halo IMBHs passing through the disk region, $d f_n/dn$ the volume fraction of regions in the Galactic disk which is occupied by molecular clouds of density n per unit cloud density, and $d f_v/dv$ the velocity distribution function of IMBHs. In this equation, v is a function of n and L_X as determined by eq. (1).

Assuming the Navarro, Frenk, & White (1997) density profile with parameters given in Klypin, Zhao, & Somerville (2002), we estimate the dark matter mass included within 10 kpc radius is about 4% of M_{halo} ¹. Then f_{disk} can be estimated as $\sim 0.04 \times (3/4) 2H_g/(10 \text{ kpc}) \sim 4.5 \times 10^{-4}$, where $H_g = 75 \text{ pc}$ (Sanders, Solomon, & Scoville 1984) is the disk scale height of molecular gas. The velocity of ULXs must be much smaller than the dispersion of the Maxwell distribution, $\sigma_{\text{BH}} = V_c/\sqrt{2} = 160 \text{ km/s}$, and hence $d f_v/dv \sim (2/\pi)^{1/2} v^2/\sigma_{\text{BH}}^3$.

Following Agol & Kamionkowski (2002), we model f_n as $d f_n/dn \propto n^{-\beta}$ with $\beta = 2.8$, between $n_{\text{min}} = 10^2$ and $n_{\text{max}} = 10^5 \text{ cm}^{-3}$. Normalization is given as:

$$\frac{d f_n}{dn} = \frac{(\beta-2) \langle \Sigma_{\text{MC}} \rangle}{2\mu m_p n_{\text{min}}^2 H_g} \left(\frac{n}{n_{\text{min}}} \right)^{-\beta}, \quad (3)$$

where $\langle \Sigma_{\text{MC}} \rangle$ is the mean surface mass density of molecular gas in the Galactic disk, which is averaged for IMBHs passing through the disk as:

$$\langle \Sigma_{\text{MC}} \rangle = \frac{\int 2\pi r \Sigma_{\text{MC}}(r) \rho_{\text{DM}} dr}{\int 2\pi r \rho_{\text{DM}} dr}, \quad (4)$$

where r is the galactocentric radius and ρ_{DM} the dark halo density. By using the data of Sanders et al. (1984) for $\Sigma_{\text{MC}}(r)$, we found $\langle \Sigma_{\text{MC}} \rangle = 29 M_\odot \text{ pc}^{-2}$.

We cannot take all value of n between n_{min} and n_{max} , because $\tilde{v}$ is limited as $\tilde{v} > (\sigma_{\text{MC}}^2 + c_s^2)^{1/2}$ and n must be larger than

$$n'_{\text{min}} \equiv 2.5 \times 10^3 L_{39} \eta_{-1}^{-1} M_2^{-2} \left[\frac{(\sigma_{\text{MC}}^2 + c_s^2)^{1/2}}{(10 \text{ km s}^{-1})} \right]^3 \text{ cm}^{-3} \quad (5)$$

for $v(n, L_X)$ to have a solution, where $L_{39} \equiv L_X/(10^{39} \text{ ergs s}^{-1})$. For n greater than n'_{min} ,

$$v(n, L_X) \sim 3.4 \eta_{-1}^{1/3} M_2^{2/3} n_2^{1/3} L_{39}^{-1/3} \text{ km s}^{-1}, \quad (6)$$

and hence we find

$$\frac{\partial^2 N_{\text{ULX}}}{\partial n \partial L_X} \propto n^{1-\beta} L_X^{-2}. \quad (7)$$

Therefore integrating over n between $\tilde{n}_{\text{min}}$ and n_{max} , where $\tilde{n}_{\text{min}} \equiv \max(n_{\text{min}}, n'_{\text{min}})$, we find the expected total number of

¹ The infall of baryonic gas into the center of a dark halo leads to adiabatic compression and a subsequent enhancement of dark matter density including halo IMBHs, which will be discussed in §7.

ULX as:

$$N_{\text{ULX}}(> L_X) \sim \frac{dN}{dL_X} L_X \quad (8)$$

$$\sim N_{\text{IMBH}} f_{\text{disk}} \frac{\tilde{n}_{\min}}{\beta-2} \frac{d f_n(\tilde{n}_{\min})}{d n} f_v[< v(\tilde{n}_{\min}, L_X)] \quad (9)$$

$$= 1.1 \times 10^{-2} L_{39}^{-1} \eta_{-1} M_2 \left(\frac{\tilde{n}_{\min}}{n_{\min}} \right)^{2-\beta}. \quad (10)$$

If $(\sigma_{\text{MC}}^2 + c_s^2)^{1/2} \sim 10 \text{ km s}^{-1}$, $n'_{\min}$ is larger than $n_{\min}$ and the number is reduced to $N_{\text{ULX}} \sim 8.6 \times 10^{-4}$ for the $M_{\text{BH}} = 100 M_\odot$ case, but $n'_{\min} < n_{\min}$ and no reduction of N_{ULX} for $M_{\text{BH}} = 10^3 M_\odot$.

There is no ULX in our Galaxy, and the number is known to be proportional to star formation rate (SFR) of galaxies (Grimm, Gilfanov, & Sunyaev 2003). Under a reasonable assumption that the amount of molecular gas clouds in a galaxy is proportional to SFR of the galaxy, the number of ULX should also scale with SFR². Then f_n of a starburst galaxy like the Antennae is 30 times larger than that of our Galaxy. The number of ULXs in such a galaxy is then estimated to be ~ 3.3 for $M_{\text{BH}} = 10^3 M_\odot$, while in the Antennae ~ 10 ULXs are observed above $10^{39} \text{ ergs s}^{-1}$ (Zezas & Fabbiano 2002). This is an encouraging result; it should be noted that the observed ULXs probably include ordinary X-ray binaries in the luminosity range of $\sim (1-3) \times 10^{39} \text{ ergs s}^{-1}$. In the following sections, we discuss two more physical processes; one (gas dynamical friction) may enhance the ULX number and the other (radiative efficiency of accretion) is important to reproduce the observed luminosity function of X-ray sources.

3. THE EFFECT OF GAS DYNAMICAL FRICTION

The expected number of ULXs is sensitively dependent on the relative velocity v , since $L_X \propto v^{-3}$ and $f_v(< v) \propto v^3$. Therefore, if there is a process that reduces the speed of IMBHs in molecular clouds, it would significantly enhance the expected ULX number. Dynamical friction is a possible effect, and considering that the star formation efficiency in molecular clouds is generally low (stellar to gas mass ratio $M_{\text{star}}/M_{\text{MC}} \sim 0.01-0.1$, Myers et al. 1986; Larson 1988), the dynamical friction by gas will be more efficient than by stars. The gas dynamical friction time scale is given by (Ostriker 1999)

$$t_{\text{df}} \equiv \frac{v}{\dot{v}} = \frac{M_{\text{BH}} v}{F_{\text{df}}} = \frac{v^3}{4\pi \xi G^2 M_{\text{BH}} \rho \ln(r_{\text{max}}/r_{\text{min}})} \quad (11)$$

$$= 8.5 \times 10^8 \xi^{-1} v_1^3 M_2^{-1} n_2^{-1} \text{ yr}, \quad (12)$$

where F_{df} is the friction force, $v_1 \equiv v/(10 \text{ km s}^{-1})$, r_{max} and r_{min} are the sizes of surrounding medium and the object receiving the force, respectively, and ξ is a parameter of order unity to take account of the uncertainty in the formula. Here we took $r_{\text{max}} = 20 \text{ pc}$ and $r_{\text{min}} = 10^3 \text{ km}$ ($\sim$ Schwarzschild radius³) for the Coulomb logarithm, $\ln(r_{\text{max}}/r_{\text{min}}) \sim 27$.

This should be compared to the crossing time of molecular clouds,

$$t_{\text{cross}} \sim \frac{L_{\text{MC}}}{v} = 2.4 \times 10^6 n_2^{-0.9} v_1^{-1} \text{ yr}, \quad (13)$$

² When the morphology of a starburst galaxy is greatly different from that of our Galaxy, a simple scaling of the ULX number with SFR may not be exactly correct. However, it is difficult to quantitatively discuss this effect and it is beyond the scope of the paper.

³ Gas within the accretion radius $r_A \sim GM_{\text{BH}}/v^2 \sim 10^{16} \text{ cm}$ is captured by the IMBH, and momentum loss by the accretion leads to a similar formula that is different only by numerical factors.

where the size of typical molecular clouds is given as $L_{\text{MC}} \sim 24.7 n_2^{-0.9} \text{ pc}$ (Larson 1981). The ratio

$$\frac{t_{\text{df}}}{t_{\text{cross}}} = 350 \xi^{-1} v_1^4 n_2^{-0.1} M_2^{-1} \quad (14)$$

becomes of order unity for $M_{\text{BH}} \sim 10^3 M_\odot$ and $v \lesssim 5 \text{ km s}^{-1}$. The dependence on v and n indicates that this is achieved only for objects having the highest accretion rate, i.e., ULXs. Therefore this process could boost up the ULX luminosity particularly at the brightest end of the ULX luminosity function.

4. RADIATIVE EFFICIENCY AND LUMINOSITY FUNCTION SLOPE

So far we assumed that the X-ray luminosity is proportional to the Bondi accretion rate, i.e., a constant radiative efficiency, η . However, in this case we have a serious problem about the ULX luminosity function. Since $L_X \propto \dot{m} \propto v^{-3}$, while $f_v(< v) \propto v^3$, a robust prediction is that the luminosity function should scale as $dN/dL \propto L^{-\alpha}$ with $\alpha = 2$. On the other hand, the observed luminosity function of ULX in star forming galaxies is smoothly connected⁴ to that of high mass X-ray binaries (HMXBs), with a slope of $\alpha = 1.6$ in a wide luminosity range of $L_X \sim 10^{36}-10^{39} \text{ ergs s}^{-1}$ (Grimm et al. 2003). Therefore, even if we successfully explained the observed number of ULXs, it would overproduce the observable number of low-luminosity X-ray sources by the same IMBH population.

This problem can be solved if we take into account the change of accretion mode at low accretion rate compared with the Eddington rate; recent studies established that accretion disks change their state from standard thin disks to radiatively inefficient accretion flows (RIAFs), at luminosity below $\sim 10\%$ of the Eddington luminosity (Kato, Fukue, & Mineshige 1998; Narayan 2004). In the RIAF regime, the luminosity scales roughly $L \propto \dot{m}^2$, and hence the luminosity function slope of ULXs becomes $\alpha = 1.5$, avoiding the overproduction problem of low luminosity sources.

5. MODELING THE LUMINOSITY FUNCTION

Here we calculate the ULX luminosity function by integrating numerically over n , v , and the location of a BH in the Galaxy. The calculation is based on the formulation of Agol & Kamionkowski (2002), but we also include the following two effects discussed above. (i) To make the luminosity proportional to $\dot{m}^2$ in the RIAF regime, the radiative efficiency is reduced as $\eta = \dot{m}/(0.1 \dot{m}_{\text{Edd}}) \eta_{\text{Edd}}$ when $\dot{m} < 0.1 \dot{m}_{\text{Edd}}$, while $\eta = \eta_{\text{Edd}}$ for $\dot{m} > 0.1 \dot{m}_{\text{Edd}}$, where $\dot{m}_{\text{Edd}} \equiv L_{\text{Edd}}/(c^2 \eta_{\text{Edd}})$, L_{Edd} the Eddington luminosity, and η_{Edd} is a constant parameter to determine the efficiency around the Eddington accretion rate. (ii) The relation $v(n, L_X)$ is replaced by $v'(n, L_X)$ to take into account the dynamical friction, where v' and v ($v' > v$) are the velocity before and after an IMBH passes through a molecular cloud having the size L_{MC} (see §3), whose relation is determined by solving the equation of motion with the dynamical friction. Since the initial mass function (IMF) of Pop III stars is highly uncertain, we simply calculate for a δ -function IMF with $M_{\text{BH}} = 10^2$ and $10^3 M_\odot$.

⁴ This fact is often used to argue that ULXs are physically an extension of normal HMXBs (e.g., Gilfanov 2004), but the latest work on ULX luminosity function indicates that ULXs with $L_X \gtrsim 10^{40} \text{ ergs s}^{-1}$ might be a different population from lower luminosity ones (Liu, Bregman, & Irwin 2005). In our scenario, the similarity between the number of IMBH-ULXs and extrapolation of the HMXB luminosity function must be a coincidence.

The results are shown in Fig. 1. We scale up the luminosity function of the Galaxy to that for a galaxy with $\text{SFR}_{>5} = 50 M_\odot \text{ yr}^{-1}$, to compare with the observed data of the universal luminosity function constructed by X-ray sources in different galaxies and normalized by SFR, where $\text{SFR}_{>5}$ is SFR for stars heavier than $5 M_\odot$ and $\text{SFR}_{>5} = 0.25 M_\odot \text{ yr}^{-1}$ for our Galaxy (Grimm et al. 2003). Considering the uncertainties, we show eight calculations with $\xi = 1$ or 10 , $c_s = 0.3 \text{ km s}^{-1}$ or 10 km s^{-1} , and $\eta_{\text{Edd}} = 0.1$ or 1 . It should be noted that the parameter η_{Edd} also includes the uncertainty in the Bondi-Hoyle accretion formula. Still, the case of $\eta_{\text{Edd}} = 1$ is rather unlikely, but we take this value as a maximally possible one. The choice of $\xi = 10$ may also be extreme, but we took this value to show the effect of dynamical friction clearly. There are two breaks in the model curves: one corresponding to the transition luminosity between the two accretion modes (the standard thin disk or RIAF) at $L_X = 0.1 L_{\text{Edd}}$, and another corresponding to the luminosity above which $\eta'_{\text{min}} > n_{\text{min}}$. As expected, the dynamical friction effect boosts up the luminosity function at the brightest luminosity range.

These results show that at least a part of the brightest ULXs ($L_X \gtrsim 10^{40} \text{ ergs s}^{-1}$) can be explained by our scenario, without overproducing low luminosity sources in the range of $L_X \lesssim 10^{39} \text{ ergs s}^{-1}$, where normal HMXBs are dominant. (Therefore we do not have to explain all of the low luminosity sources.) Especially, with $M_{\text{BH}} = 10^3 M_\odot$, the number of the most luminous ULXs ($L_X \gtrsim 10^{40} \text{ erg/s}$) can be explained with a modest value of $\eta_{\text{Edd}} = 0.1$, favoring a larger IMBH mass. The enhancement of IMBH density around galactic disks by baryon compression may further increase the number of ULXs, possibly allowing a lower IMBH mass (see §7). If the radiative feedback and ionization of ambient gas are significant, c_s should be raised to $\sim 10 \text{ km s}^{-1}$ and the number of ULX would be significantly reduced, especially in the case of $M_{\text{BH}} = 100 M_\odot$. However, in the next section we will argue that this effect is not necessarily strong.

6. ON THE RADIATIVE FEEDBACK

The ULX luminosity is sufficient to ionize the surrounding medium; the Strömgren radius is

$$r_S \sim 7.0 \times 10^{18} n_2^{-2/3} \epsilon_{-1}^{1/3} L_{39}^{1/3} \text{ cm}, \quad (15)$$

where ϵ is the fraction of ionizing luminosity above 13.6 eV compared with L_X , and $\epsilon_{-1} = \epsilon/0.1$. This is in most cases larger than the Bondi accretion radius,

$$r_A \sim \frac{GM_{\text{BH}}}{\tilde{v}^2} = 1.3 \times 10^{16} M_2 \tilde{v}_1^{-2} \text{ cm}. \quad (16)$$

Therefore, when an ULX ionizes the surrounding medium, the Bondi accretion rate should reduce, because of the increased c_s and expansion of gas driven by increased pressure. However, the gas that has already accreted to form an accretion disk will serve as the energy source of the ULX phenomenon for some time after the ionization. Based on the interstellar density gradient and angular momentum conservation, the interstellar gas captured by an IMBH should form a rotationally supported disk at a radius $r_d \sim 7.5 \times 10^{13} M_2^{5/3} v_1^{-10/3} \text{ cm}$ (Agol & Kamionkowski 2002), which is much smaller than the Bondi accretion radius, r_A . Assuming that the further accretion proceeds by α -viscosity in the standard thin disk, accretion rate for ULX-level luminosity is not achieved until a sufficient amount of mass is accumulated and the surface mass density Σ at $r \sim r_d$ reaches that of the standard disk for the mass accretion rate of ULXs, $\dot{m}$. It would

take a time of $\sim 2\pi \Sigma r_d^2 / \dot{m}$. Then ULX activity will last for a time scale of the disk mass consumption, $2\pi \Sigma r_d^2 / \dot{m}$, even after the ULX ionizes the surrounding medium.

Therefore both the time scale of turning on of ULXs and duration of the ULX phenomenon are given by the accretion time scale of the standard disk,

$$t_{\text{sd}} \sim \frac{2\pi \Sigma r_d^2}{\dot{m}} \quad (17)$$

$$\sim 2.7 \times 10^5 \alpha_{-1}^{-4/5} M_2^{-4/9} L_{39}^{49/45} \eta_{-1}^{-49/45} n_2^{-25/18} \text{ yr}. \quad (18)$$

(The outer region of the three regions in the standard disk theory applies at $r \sim r_d$.) It is interesting that this time scale for an ULX with $L_X \sim 10^{40} \text{ ergs s}^{-1}$ is similar to the crossing time scale $t_{\text{cross}} \sim 10^6 \text{ yr}$ of a $n \sim 10^2 \text{ cm}^{-3}$ cloud (§3). This means that the duty cycle as an ULX could effectively be of order unity for an IMBH passing through a molecular cloud, even if the radiative feedback is significant.

7. DISCUSSION AND CONCLUSIONS

We conclude that the observed number of ULXs can be explained by halo population IMBHs passing through dense molecular clouds, which were produced as Pop III star remnants, if the Pop III star formation was very efficient as recently indicated by the unaccounted excess of the cosmic near-infrared background radiation. Luminosity function was calculated, and the ULXs of our scenario contribute to the observed luminosity function especially at the largest luminosity range, without overproducing sources in low-luminosity range where stellar-mass black holes are the dominant contribution. The ULX phenomena could last for 10^{5-6} yr and the duty cycle in a molecular cloud could be of order unity.

We considered molecular clouds with a density range of 10^2 – 10^5 cm^{-3} , but rather low-density clouds of $n \sim 10^2 \text{ cm}^{-3}$ have the largest contribution to the expected ULX number. It is consistent with observations of ULX locations; many ULXs are found in star forming regions but in outskirts of large concentrations of gas (Zezas et al. 2002).

Large ionized, expanding optical nebulae with a size of $\gtrsim 100 \text{ pc}$ are often associated with ULXs, with estimated total expansion energy of $\sim 10^{52} \text{ erg}$ and age of $\sim 10^6 \text{ yr}$ (Pakull & Mirioni 2002; Mushotzky 2004). Our estimate of the ULX duration is consistent with this age, and the total energy emitted by an ULX with $L_X \sim 10^{40} \text{ ergs s}^{-1}$ could be as large as $\sim 3 \times 10^{53} \text{ erg}$. A part of this energy may be emitted in UV and soft X-ray bands to ionize and expand the ambient medium, giving an explanation for the observed optical nebulae.

The estimated ULX duration is not very different from the Eddington time scale, $t_{\text{Edd}} = M_{\text{BH}} / (L_{\text{Edd}} / \eta c^2) = 4.4 \times 10^7 \eta_{-1} \text{ yr}$. Therefore, some ULXs may have a chance to accrete gas for a time scale comparable to or longer than t_{Edd} , by passing through a large molecular cloud or many molecular clouds. Such IMBHs will increase their mass considerably. In fact, the IMBH mass used in our calculation is not necessarily that of the Pop III remnants. The mass function of IMBHs in the local universe may have been altered from the original one at the formation epoch, by merging or gas accretion in the hierarchical structure formation. Such calculations have been performed by Islam, Taylor, & Silk (2004) and Volonteri & Perna (2005). According to these calculations, typical mass of free-floating IMBHs may be increased up to $\sim 10^4 M_\odot$ depending on model parameters, even if the initial mass of Pop III remnant BHs is less than $1000 M_\odot$. The very large BH

mass suggested for the extremely bright ULX in M82 (Matsumoto et al. 2001; Kaaret et al. 2001) may be achieved by such processes.

We used the popular NFW density profile for the Galactic halo, but the dark matter density could be significantly altered and increased around the region of the Galactic disk, by adiabatic compression during the infall of baryonic gas to form stars. The number of ULXs would then be increased, because more IMBHs exist around the Galactic disk and the weighted mean surface density of molecular gas is also increased. According to the model by Klypin et al. (2002), we estimate the boost factor of the ULX number as ~ 3 . On the other hand, the same authors also shown that a modest exchange of angular momentum between baryonic and dark matter (through e.g., a rotating bar-like bulge) may result in almost no enhancement, or even decrease of the dark matter density. This should be taken into consideration as an uncertainty in the prediction made here, though the main conclusions are unlikely to be changed.

Finally we give a list of the predictions for ULXs produced by the scenario proposed here. Since ULXs are isolated IMBHs, optical counterparts (donor stars) will not be

found. Future observations should confirm the trend that ULXs are preferentially found in star forming regions, but at outskirts of dense molecular clouds where the density is about $\sim 100 \text{ cm}^{-3}$. Expanding nebulae with kinetic energy of $\sim 10^{52} \text{ erg}$ should often be associated, *without* evidence for a past hypernova or multiple supernovae. The predicted slope of the IMBH luminosity function is $\alpha = 1.5$ at low luminosity range, which is similar to that of HMXBs. Therefore, in a wide range of luminosity, we expect that a small fraction of low-luminosity X-ray sources are the same population with IMBH-ULXs, showing RIAF spectra without companion stars. It is interesting to search for such a population in the Galaxy or nearby galaxies. The excess of CNIB must be confirmed with greater significance. The high density of free-floating IMBHs in galactic haloes may be detected by future gravitational lensing experiments (e.g., Totani 2003).

We would like to thank the anonymous referee for useful comments. This work has partially been supported by the Grant-in-Aid for the 21st Century COE "Center for Diversity and Universality in Physics" from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan.

REFERENCES

- Agol, E. & Kamionkowski, M. 2002, MNRAS, 334, 553
 Begelman, M.C. 2002, ApJ, 568, L97
 Cambr  s, L., Reach, W.T., Beichman, C.A. & Jarrett T.H., 2001, ApJ, 555, 563
 Colbert, E.J.M. & Miller, M.C. 2004, to appear in the proceedings of the Tenth Marcel Grossmann Meeting on General Relativity, ed. M. Novello, et al., World Scientific, Singapore (astro-ph/0402677)
 Cropper, M., Soria, R., Mushotzky, R.F., Wu, K., Markwardt, C.B. & Pakull, M. 2004, MNRAS, 349, 39
 Gilfanov, M. 2004, Progress of Theoretical Physics, in press. (Proceedings of the workshop "Stellar-Mass, Intermediate-Mass, and Supermassive Black Holes", Eds: K.Makishima and S.Mineshige.) (astro-ph/0403552)
 Fryer, C.L. & Kalogera, V. 2001, ApJ, 554, 548
 Fujita, Y., Inoue, S., Nakamura, T., Manmoto, T. & Nakamura, K.E. 1998, ApJ, 495, L85
 Grimm, H.J., Gilfanov, M. & Sunyaev R. 2003, MNRAS, 339, 793
 Islam, R.R., Taylor, J.E. & Silk, J. 2004, ApJ, 354, 443
 Kaaret, P., Prestwich, A.H., Zezas, A., Murray, S.S., Kim, D.-W., Kilgard, R.E., Schlegel, E.M. & Ward, M.J. 2001, MNRAS, 321, L29
 Kato, S., Fukue, J. & Mineshige, S. 1998, Black-Hole Accretion Disks (Kyoto:Kyoto Univ. Press)
 King, A.R. 2004, ApJ, 347, L18
 King, A.R., Davies, M.B., Ward, M.J., Fabbiano, G. & Elvis, M. 2001, ApJ, 552, L109
 Klypin, A., Zhao, H. & Somerville, R.S. 2002, ApJ, 573, 597
 Krolik, J.H. 2004, ApJ, 615, 383
 Larson, R.B. 1981, MNRAS, 194, 809
 Larson, R. B. 1988, in 'Galactic and Extragalactic Star Formation', eds. R.E. Pudritz and M. Fich (Kluwer Academic Publishers)
 Liu, J.-F., Bregman, J.N., & Irwin, J. 2005, a preprint submitted to ApJ, (astro-ph/0501312)
 Matsumoto, H., Tsuru, T.G., Koyama, K., Awaki, H., Canizares, C.R., Kawai, N., Matsushita, S. & Kawabe, R. 2001, ApJ, 547, L25
 Matsumoto, T. et al. 2004, to appear in ApJ (astro-ph/0411593)
 Madau, P. & Rees, M.J. 2001, ApJ, 551, L27
 Madau, P. & Silk, J. 2005, MNRAS, in press (astro-ph/0502304)
 Makishima, K. et al. 2000, ApJ, 535, 632
 Miller, J. M., Fabian, A. C., & Miller, M. C. 2004, ApJ, 614, L117
 Murali, C., Arras, P. & Wasserman, I. 2000, MNRAS, 313, 87
 Mushotzky, R. 2004, astro-ph/0411040
 Myers, P.C., Dame, T.M., Thaddeus, P., Cohen, R.S., Silverberg, R.F., Dwek, E. & Hauser, M.G. 1986, ApJ, 301, 398
 Narayan, R. 2004, To appear in "From X-ray Binaries to Quasars: Black Hole Accretion on All Mass Scales" ed. T. Maccarone, R. Fender, L. Ho (Kluwer), astro-ph/0411385
 Navarro, J.F., Frenk, C.S. & White, S.D.M. 1997, ApJ, 490, 493
 Ostriker, E.C. 1999, ApJ, 513, 252
 Pakull, M.W. & Mirioni, L. 2002, in 'New Visions of the X-ray Universe in the XMM-Newton and Chandra Era' (astro-ph/0202488)
 Portegies Zwart, S.F. & McMillan, S.L.W. 2002, ApJ, 576, 899
 Salvaterra, R. & Ferrara, A. 2003, MNRAS, 339, 973
 Sanders, D.B., Solomon, P.M. & Scoville, N.Z. 1984, ApJ, 276, 182
 Santos, M.R., Bromm, V. & Kamionkowski, M. 2002, MNRAS, 336, 1082
 Schneider, R., Ferrara, A., Natarajan, P. & Omukai, K. 2002, ApJ, 571, 30
 Scoville, N.Z. & Sanders, D.B. 1987, in Hollenbach D.J., Thronson H.A., eds, Interstellar Processes. Reidel, Dordrecht, p. 21
 148,
 Totani, T., Yoshii, Y., Iwamuro, F., Maihara, T. & Motohara, K. 2001, ApJ, 550, L137
 Totani, T. 2003, ApJ, 586, 735
 Volonteri, M. & Perna, R. 2005, to appear in MNRAS, astro-ph/0501345
 Wambsganss, J. 2002, in 'Where's the Matter? Tracing Bright and Dark Matter with the New Generation of Large Scale Surveys', Eds. M. Treyer & L. Tresse (astro-ph/0207616)
 Wasserman, I. & Salpeter, E. E. 1994, ApJ, 433, 670
 Wright, E.L. & Reese, E.D. 2000, ApJ, 545, 43
 Yoo, J., Chanam  , J. & Gould, A. 2004, ApJ, 601, 311
 Zezas, A. & Fabbiano, G. 2002, ApJ, 577, 726
 Zezas, A., Fabbiano, G., Rots, A.H. & Murray, S.S. 2002, ApJ, 577, 710




FIG. 1.— *Left:* The X-ray luminosity function of free-floating IMBHs ($M_{\text{BH}} = 100 M_{\odot}$) passing through dense molecular clouds, normalized to $\text{SFR}_{>5} = 50 M_{\odot} \text{ yr}^{-1}$. For each line marking, there are two model curves: the upper one for $\eta_{\text{Edd}} = 1$ and the lower one for $\eta_{\text{Edd}} = 0.1$. The sound speed is set as $c_s = 0.3$ and 10 km s^{-1} for each thick and thin curve, respectively. The dynamical friction parameter is set to be $\xi = 1$ and 10 for each solid and dashed curve, respectively. The data points show the observed universal X-ray luminosity function from Grimm et al. (2003). *Right:* Same as the left panel, but for $M_{\text{BH}} = 10^3 M_{\odot}$.